

RPA GREEN'S FUNCTIONS OF THE ANISOTROPIC HEISENBERG MODEL

Andre Johannes Stoffel and Miklós Gulácsi

Max-Planck-Institute for the Physics of Complex Systems, D-01187 Dresden, Germany

Nonlinear Physics Centre, Australian National University, Canberra, ACT 0200, Australia

November 10, 2009

We solve in random-phase approximation the anisotropic Heisenberg model, including nearest and next-nearest neighbour interactions by calculating all Green's functions and pair correlation functions in a cumulant decoupling scheme. The general exposition is pedagogic in tone and is intended to be accessible to any graduate student or physicist who is not an expert in the field.

I. INTRODUCTION

Recently we analysed the properties of an anisotropic Heisenberg model in an external longitudinal field on a bcc lattice, with a particular application to supersolids [2, 3, 4]. However, detailed derivation of the Green's functions and correlation functions have not yet been published, as such we fill this gap hereafter. For the interested reader we present in Appendix A the connection between the model analysed hereafter and ^4He .

We work with the standard anisotropic Heisenberg model, defined by the Hamiltonian:

$$H = h^z \sum_i S_i^z + \sum_{ij} J_{ij}^{\parallel} S_i^z S_j^z + \sum_{ij} J_{ij}^{\perp} (S_i^x S_j^x + S_i^y S_j^y) \quad (1)$$

on a bcc lattice shown in Figure (1). It is known that the Hamiltonian defined in Equation (1) gives four magnetic phases: ferromagnetic, canted ferromagnetic, canted anti-ferromagnetic and anti-ferromagnetic phases which we abbreviate by FE, CFE, CAF and AF, see Table (I). The order parameters, m_1 for off-diagonal long range order and m_2 for diagonal long range order in the magnetic system are defined by:

$$\begin{aligned} m_1 &= \langle S_A^x \rangle + \langle S_B^x \rangle \\ m_2 &= \langle S_A^z \rangle - \langle S_B^z \rangle \end{aligned} \quad (2)$$

In the following we will use these order parameters to identify the phases within the random-phase approximation.




FIG. 1: The bcc lattice consists of two interpenetrating sc sub-lattices, i.e., sub-lattice A and sub-lattice B. For simplicity we only have drawn the two dimensional case.

The Heisenberg model has been studied thoroughly although there exists only a classical mean-field solution for the canted ferromagnetic and anti-ferromagnetic phases [5, 6]. Hence, our interest is in obtaining a better than mean-field solution of these canted phases. We used [2, 3, 4] the equation of motion technique for the time-temperature-dependent Green's function [7], $G_{ij}^{xy}(t)_{Ret/Adv} = \mp i\theta(\pm t)\langle[S_i^x(t), S_j^y]\rangle$, for all four phases, which we solve in random-phase approximation (RPA). In order to maintain pair correlations as accurate as possible, we have chosen a cumulant decoupling scheme to approximate higher order Green's functions beyond the random-phase approximation. This is the first time that a thorough cumulant random-phase approximation many-body calculation has been solved for the canted phases and as such it represents a vital step in the understanding the behaviour of the three dimensional anisotropic Heisenberg model in the presence of an external magnetic field. Details of the Green's functions are given in the next Chapter.

By using these Green's functions we calculate exactly all the pair correlations, also given in the next Chapter, where we show that this decoupling scheme couples six Green's functions, one for each spin component in x , y and z direction on the two sub-lattices respectively, to a set of six equations. Due to the enormous number of terms (1024 in total) within these Green's functions we have decided to reproduce their exact form only at the end. Using these Green's functions it can be shown or alternatively argued that the Goldstone theorem of gapless modes imposes an additional condition [8] on the mean fields of the

Spin Configuration	Magnetic Phase	ODLRO	DLRO
$\uparrow\uparrow$	FE	No	No
$\nearrow\nearrow$	CFE	Yes	No
$\nearrow\nwarrow$	CAF	Yes	Yes
$\uparrow\downarrow$	AF	No	Yes

TABLE I: Possible magnetic phases of the anisotropic Heisenberg model. All phases are defined by their long range order, i.e., off-diagonal long-range order (ODLRO) and diagonal long-range order (DLRO).

canted phases, reducing the number of order-parameters by two. As this condition does not apply to the ferromagnetic and anti-ferromagnetic phases, their Green's functions are structurally different.

For completeness we also re-derive the classical mean-field Green's functions in Appendix B. While in Appendix C we establish a link between the classical mean-field and the random-phase approximations.

II. GREEN'S FUNCTIONS

The most versatile and successful method of solving many-body problems involve the concept of Green's function. The Matsubara formalism [7] for imaginary time Green's functions at finite temperature is commonly used to analyze fermionic and bosonic many-body systems. However, in spin systems a different type of Green's function is primarily applied as this makes it easier to deal with the algebraic properties of spin operators. The retarded and advanced Tyablikov [9, 10] commutator Green's function at finite temperature defined

in real time are:

$$\begin{aligned} G_{ijRet}^{\mu\nu}(t) &= -i\theta(t)\langle[S_i^\mu(t), S_j^\nu]\rangle \\ G_{ijAdv}^{\mu\nu}(t) &= i\theta(-t)\langle[S_i^\mu(t), S_j^\nu]\rangle \end{aligned} \quad (3)$$

The average $\langle \rangle$ involves the usual quantum mechanical as well as thermal averages and μ and ν are elements of $\{x, y, z\}$ and i and j denote the lattices sites. There is no Wick's theorem for spin systems and therefore no perturbative approaches by means of Feynman diagrams are available. However this is not necessarily a disadvantage as the method of the equation of motion is equally powerful in many cases and often much more compact. The basic idea of this method is to find a linear differential equation for the Green's function. Therefore we can differentiate the xy -Green's function with respect to time:

$$\begin{aligned} i\partial_t G_{ijRet}^{xy}(t) &= \delta(t)\langle[S_i^x, S_j^y]\rangle - i\theta(t)\langle[[S_i^x, H], S_j^y]\rangle \\ i\partial_t G_{ijAdv}^{xy}(t) &= \delta(t)\langle[S_i^x, S_j^y]\rangle + i\theta(-t)\langle[[S_i^x, H], S_j^y]\rangle \end{aligned} \quad (4)$$

Here we have used the Heisenberg equation of motion $i\partial_t S^x = [S_i^x, H]$ for the operator S^x . For the anisotropic Heisenberg model the commutator yields:

$$[S_i^x, H] = -i\mu S_i^y - 2i \sum_j J_{ij}^{\parallel} S_i^y S_j^z + 2i \sum_j J_{ij}^{\top} S_i^z S_j^y \quad (5)$$

Consequently the right hand side of Equation (4) involves higher, third order Green's functions. In principle those third order functions can be expressed in terms of even higher order. This procedure will go on indefinitely yielding a series of infinite order. The central idea is to introduce a suitable approximation which truncates this series. Here, in order to keep the complexity to a manageable level we split up the three operator correlation functions.

The cumulant decoupling [11] is based on the assumption that the last term of the following equality is negligible:

$$\begin{aligned} \langle \hat{A}\hat{B}\hat{C} \rangle &= \\ &\langle \hat{A} \rangle \langle \hat{B}\hat{C} \rangle + \langle \hat{B} \rangle \langle \hat{A}\hat{C} \rangle \\ &+ \langle \hat{C} \rangle \langle \hat{A}\hat{B} \rangle - 2\langle \hat{A} \rangle \langle \hat{B} \rangle \langle \hat{C} \rangle \\ &+ \langle (\hat{A} - \langle \hat{A} \rangle)(\hat{B} - \langle \hat{B} \rangle)(\hat{C} - \langle \hat{C} \rangle) \rangle \end{aligned} \quad (6)$$

This approximation is justified if the quantum fluctuations are small and do not deviate far from their mean field values, which is the case in three dimensions [12]. As a consequence the third order correlation functions split into product terms of a second order correlation functions and the mean-field of a single operators:

$$\begin{aligned} \langle [[S_i^\alpha(t), S_k^\beta(t)], S_j^y] \rangle \rightarrow \\ \langle S_i^\alpha(t) \rangle \langle [S_k^\beta(t), S_j^y] \rangle + \langle S_k^\beta(t) \rangle \langle [S_i^\alpha(t), S_j^y] \rangle \end{aligned} \quad (7)$$

Finally we obtain a differential equation which only involves second order Green's functions.

$$\begin{aligned} i\partial_t G_{ij}^{xy}(t) = \\ i\delta(t)\delta_{ij}\langle S_i^z \rangle - i\mu G_{ij}^{yy}(t) \\ -2i \sum_l J_{il}^\parallel (\langle S_i^y(t) \rangle G_{lj}^{zy}(t) + \langle S_l^z(t) \rangle G_{ij}^{yy}(t)) \\ +2i \sum_l J_{il}^\top (\langle S_i^z(t) \rangle G_{lj}^{yy}(t) + \langle S_l^y(t) \rangle G_{ij}^{zy}(t)) \end{aligned} \quad (8)$$

Here we have dropped the subscripts for the retarded and the advances Green's functions since the equations equally hold for both. In the next Chapter we will see that the advanced and the retarded Green's functions are actually represented by the same Fourier transform; the difference comes from the path along which the inverse Fourier integral is carried out. Similar relations also exist for the time derivatives of $G_{ij}^{yy}(t)$ and $G_{ij}^{zy}(t)$. Together these equations form a closed set of linear differential equations. Usually linear differential equations are readily solved by Fourier Transformation. The time-frequency Fourier and inverse Fourier transforms are given by:

$$G_{ij}^{xy}(t) = \int d\omega e^{-i\omega t} G_{ij}^{xy}(\omega) \quad (9)$$

$$G_{ij}^{xy}(\omega) = \frac{1}{2\pi} \int dt e^{i\omega t} G_{ij}^{xy}(t) \quad (10)$$

One might think that due to broken translational symmetry of the canted anti-ferromagnetic and the anti-ferromagnetic phases, featuring diagonal long-range order, a spacial Fourier transform might not be applicable. Actually, the canted anti-ferromagnetic and anti-ferromagnetic states do exhibit discrete translational symmetry, namely they are invariant under $r_{ij} \rightarrow r_{ij} + n_1 a_1 + n_2 a_2 + n_3 a_3$, where a_1 , a_2 and a_3 are the basic lattice vectors of the sc sub-lattice and $n_1, n_2, n_3 \in Z$. This translation maps each sub-lattice onto itself and

the number of equation doubles as we have to treat each sub-lattice separately. The Fourier transform into k-space is defined by:

$$\begin{aligned} G_{ij}^{xy}(\omega) &= \int d^3k e^{ikR_{ij}} G^{xy}(k, \omega) \\ G^{xy}(k, \omega) &= \frac{1}{(2\pi)^3 n} \sum_j e^{-ikR_{ij}} G_{ij}^{xy}(\omega) \end{aligned} \quad (11)$$

After successively carrying out time and space Fourier transforms we derive a set of six algebraic equations, determining six Green's functions that represent the xy-,yy-, and zy-spin correlations on each sub-lattice. The detailed calculation is carried out at the end of this Chapter. In matrix form this set of equations reads:

$$M \cdot \gamma = v \quad (12)$$

where

$$M = \begin{pmatrix} i\omega & 0 & M_{13} & M_{14} & M_{15} & M_{16} \\ 0 & i\omega & M_{23} & M_{24} & M_{25} & M_{26} \\ -M_{13} & -M_{14} & i\omega & 0 & M_{35} & M_{36} \\ -M_{23} & -M_{24} & 0 & i\omega & M_{45} & M_{46} \\ M_{51} & M_{52} & M_{53} & M_{54} & i\omega & 0 \\ M_{61} & M_{62} & M_{63} & M_{64} & 0 & i\omega \end{pmatrix} \quad (13)$$

and

$$\gamma = \begin{pmatrix} G_{a_k}^{xy}(\omega) \\ G_{b_k}^{xy}(\omega) \\ G_{a_k}^{yy}(\omega) \\ G_{b_k}^{yy}(\omega) \\ G_{a_k}^{zy}(\omega) \\ G_{b_k}^{zy}(\omega) \end{pmatrix} \quad (14)$$

as well as

$$v = \frac{1}{(2\pi)^4} \begin{pmatrix} \langle S_A^z \rangle \\ \langle S_B^z \rangle \\ 0 \\ 0 \\ \langle -S_A^x \rangle \\ \langle -S_B^x \rangle \end{pmatrix} \quad (15)$$

The components of the matrix M are given by:

$$\begin{aligned}
M_{13} &= 2h^z + 4\langle S_A^z \rangle (J_2^\parallel(0) - J_2^\top(k)) + 4\langle S_B^z \rangle J_1^\parallel(0) \\
M_{14} &= -4\langle S_A^z \rangle J_1^\top(k) \\
M_{23} &= -4\langle S_B^z \rangle J_1^\top(k) \\
M_{24} &= 2h^z + 4\langle S_B^z \rangle (J_2^\parallel(0) - J_2^\top(k)) + 4\langle S_A^z \rangle J_1^\parallel(0) \\
M_{15} &= 4\langle S_A^y \rangle (J_2^\parallel(k) - J_2^\top(0)) - 4\langle S_B^y \rangle J_1^\top(0) \\
M_{16} &= 4\langle S_A^y \rangle J_1^\parallel(k) \\
M_{25} &= 4\langle S_B^y \rangle J_1^\parallel(k) \\
M_{26} &= 4\langle S_B^y \rangle (J_2^\parallel(k) - J_2^\top(0)) - 4\langle S_A^y \rangle J_1^\top(0) \\
M_{35} &= -4\langle S_A^x \rangle (J_2^\parallel(k) - J_2^\top(0)) + 4\langle S_B^x \rangle J_1^\top(0) \\
M_{36} &= -4\langle S_A^x \rangle J_1^\parallel(k) \\
M_{45} &= -4\langle S_B^x \rangle J_1^\parallel(k) \\
M_{46} &= -4\langle S_B^x \rangle (J_2^\parallel(k) - J_2^\top(0)) + 4\langle S_A^x \rangle J_1^\top(0) \\
M_{51} &= 4\langle S_B^y \rangle J_1^\top(0) + 4\langle S_A^y \rangle (J_2^\top(0) - J_2^\top(k)) \\
M_{52} &= -4\langle S_A^y \rangle J_1^\top(k) \\
M_{61} &= -4\langle S_B^y \rangle J_1^\top(k) \\
M_{62} &= 4\langle S_A^y \rangle J_1^\top(0) + 4\langle S_B^y \rangle (J_2^\top(0) - J_2^\top(k)) \\
M_{53} &= -4\langle S_B^x \rangle J_1^\top(0) - 4\langle S_A^x \rangle (J_2^\top(0) - J_2^\top(k)) \\
M_{54} &= 4\langle S_A^x \rangle J_1^\top(k) \\
M_{63} &= 4\langle S_B^x \rangle J_1^\top(k) \\
M_{64} &= -4\langle S_A^x \rangle J_1^\top(0) - 4\langle S_B^x \rangle (J_2^\top(0) - J_2^\top(k))
\end{aligned} \tag{16}$$

Here the k-dependent coupling constants are defined by $J_1^\top(k) = J_1^\top \gamma_1(k)$, $J_2^\top(k) = J_2^\top \gamma_2(k)$, $J_1^\parallel(k) = J_1^\parallel \gamma_1(k)$ and $J_2^\parallel(k) = J_2^\parallel \gamma_1(k)$, where

$$\begin{aligned}
\gamma_1(\mathbf{k}) &= \frac{1}{q_1} \sum_{a_{AB}} e^{i\mathbf{k} \cdot \mathbf{a}_{AB}} \\
\gamma_2(\mathbf{k}) &= \frac{1}{q_2} \sum_{a_{AA}} e^{i\mathbf{k} \cdot \mathbf{a}_{AA}}
\end{aligned} \tag{17}$$

are the lattice generating functions. On the bcc lattice these lattice generating functions are given by

$$\begin{aligned}\gamma_1(\mathbf{k}) &= \cos\left(\frac{k_x a}{2}\right) \cos\left(\frac{k_y a}{2}\right) \cos\left(\frac{k_z a}{2}\right) \\ \gamma_2(\mathbf{k}) &= \frac{\cos(k_x a)}{3} + \frac{\cos(k_y a)}{3} + \frac{\cos(k_z a)}{3}\end{aligned}\tag{18}$$

where a is the lattice constant of a simple cubic sub-lattice. Again, the spontaneously broken U(1)-symmetry gives us the freedom to set $\langle S_A^y \rangle = \langle S_B^y \rangle = 0$. This reduces the number of non-zero matrix components of M .

It has been shown that the commutator Green's functions must not have a zero frequency pole [8]. This results directly from the fact that the commutator of to spin operators at long time distances becomes zero: $\lim_{t \rightarrow \infty} \langle [S^\mu(t) S^\nu] \rangle = 0$. In the present calculation the Green's functions do actually acquire a zero frequency pole, as given by the eigenvalues of the matrix M . In the ferromagnetic and anti-ferromagnetic phases this zero frequency pole is readily canceled out, but for the canted ferromagnetic and the canted anti-ferromagnetic phases this imposes an additional constraint:

$$\begin{aligned}h^z + 2\langle S_A^z \rangle (J_2^\parallel - J_2^\top) + 2\langle S_B^z \rangle J_1^\parallel &= 2J_1^\top \frac{\langle S_B^x \rangle}{\langle S_A^x \rangle} \langle S_A^z \rangle \\ h^z + 2\langle S_B^z \rangle (J_2^\parallel - J_2^\top) + 2\langle S_A^z \rangle J_1^\parallel &= 2J_1^\top \frac{\langle S_A^x \rangle}{\langle S_B^x \rangle} \langle S_B^z \rangle\end{aligned}\tag{19}$$

$$\tag{20}$$

Note that these two conditions are identical to the classical mean-field equations [2, 3, 4]. We use these two relations to replace the external magnetic field h_z in the matrix M . To combine terms we introduce following variables:

$$\begin{aligned}A_1 &= J_2^\top (1 - \gamma_2(k)) + J_1^\top \frac{\langle S_B^x \rangle}{\langle S_A^x \rangle} \\ A_2 &= J_2^\top (1 - \gamma_2(k)) + J_1^\top \frac{\langle S_A^x \rangle}{\langle S_B^x \rangle} \\ B_1 &= J_2^\top (1 - \gamma_2(k)) + J_1^\top \frac{\langle S_B^x \rangle}{\langle S_A^x \rangle} \\ &\quad - 4(J_2^\parallel - J_2^\top) \gamma_2(k) \langle S_A^x \rangle^2 \\ B_2 &= J_2^\top (1 - \gamma_2(k)) + J_1^\top \frac{\langle S_A^x \rangle}{\langle S_B^x \rangle} \\ &\quad - 4(J_2^\parallel - J_2^\top) \gamma_2(k) \langle S_B^x \rangle^2\end{aligned}$$

$$C = J_1^\top \gamma_1(k)$$

$$D = -4\gamma_1(k)(J_1^\top \langle S_A^z \rangle \langle S_B^z \rangle + 2J_1^\parallel \langle S_A^x \rangle \langle S_B^x \rangle) \quad (21)$$

Then the matrix M in the canted anti-ferromagnetic and canted ferromagnetic phases, where Equation (19) holds, is given by (in block-form):

$$M_c = \begin{pmatrix} M_{11}^c & M_{12}^c & M_{13}^c \\ M_{21}^c & M_{22}^c & M_{23}^c \\ M_{31}^c & M_{32}^c & M_{33}^c \end{pmatrix} \quad (22)$$

where the 2×2 -blocks are given by:

$$M_{11}^c = M_{22}^c = M_{33}^c = \begin{bmatrix} i\omega & 0 \\ 0 & i\omega \end{bmatrix}$$

$$M_{12}^c = -M_{21}^c = \begin{bmatrix} 4A_1 \langle S_A^z \rangle & -4C \langle S_A^z \rangle \\ -4C \langle S_B^z \rangle & 4A_2 \langle S_B^z \rangle \end{bmatrix}$$

$$M_{13}^c = M_{31}^c = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$M_{23}^c = \begin{bmatrix} -\frac{4A_1 \langle S_A^z \rangle^2 - B_1}{\langle S_A^x \rangle} & \frac{4C \langle S_A^z \rangle \langle S_B^z \rangle + D}{\langle S_B^x \rangle} \\ \frac{4C \langle S_A^z \rangle \langle S_B^z \rangle + D}{\langle S_A^x \rangle} & -\frac{4A_2 \langle S_B^z \rangle^2 - B_2}{\langle S_B^x \rangle} \end{bmatrix}$$

$$M_{32}^c = \begin{bmatrix} 4 - A_1 \langle S_A^x \rangle & 4C \langle S_A^x \rangle \\ 4C \langle S_B^x \rangle & -4A_2 \langle S_B^x \rangle \end{bmatrix} \quad (23)$$

In the phases that do not exhibit off-diagonal long range order, namely the ferromagnetic and anti-ferromagnetic phases, Equation (19) does not hold. However, as $\langle S_x \rangle = \langle S_y \rangle = 0$ the matrix M reduced to a matrix of 4×4 -dimensions:

$$M^{\text{nc}} = \begin{pmatrix} i\omega & 0 & M_{13} & M_{14} \\ 0 & i\omega & M_{23} & M_{24} \\ -M_{13} & -M_{14} & i\omega & 0 \\ -M_{23} & -M_{24} & 0 & i\omega \end{pmatrix} \quad (24)$$

Equation (12) readily implies that the Green's functions are given by:

$$M^{-1} \cdot v \quad (25)$$

As these Green's functions in explicit form are rather complex, we defer presenting them in their full form until the end of this Chapter. Those readers who are not interested in the details of these functions might safely skip the corresponding section of the Chapter.

The Green's functions derived above determine the state of the system and all relevant macroscopic and thermodynamic properties can be extracted from them. Nevertheless the Green's functions as derived in this random phase approximation are function not only of the external field h^z but also of the mean fields of the spins. Therefore we have to define self-consistency equation which determine those mean-fields.

III. SELF-CONSISTENCY EQUATIONS AT ZERO TEMPERATURE

The Green's functions as derived in the previous Chapter do not bear an explicit dependence on the temperature but rather depend on the temperature through the various mean-fields. Thus, upon deriving the determining self-consistency equations the temperature will be introduced explicitly into the formalism. We have decided here to separate the zero temperature and the finite temperature formalism. Even though, it would generally be easy to carry out the limiting process $T \rightarrow 0$ at any time it would cause numerical difficulties to do so at a later stage as the temperature usually appears in terms of $1/T$. Also we find it instructive to derive the zero temperature formalism separately as it provides unveiled insight into the quantum nature of the model.

Naturally there seem to be three ways to set up self-consistency equations. We could, for example calculate:

$$\begin{aligned}
-\frac{1}{2}\langle S_i^z \rangle &= \langle i[S_i^x(0), S_i^y] \rangle \\
&= G_{ij_{Adv}}^{xy}(0) - G_{ij_{Ret}}^{xy}(0) \\
&= \int_{-\infty}^{\infty} d\omega [G_{ij_{Adv}}^{xy}(\omega) - G_{ij_{Ret}}^{xy}(\omega)]
\end{aligned} \tag{26}$$

an equivalent relation holds for $\langle S_i^x \rangle$ and G_{ij}^{zy} . Here, for the sake of readability we have disregarded the sub-lattice subscripts A and B. Unfortunately, these equations are not suitable to calculate the mean-fields self-consistently but rather lead to an identity, giving identically $\frac{i}{2}\langle S^z \rangle$ on both sides. That is due to the structure of the equation of motion, and the $\langle S_i^z \rangle$ are exactly those ones contained in the right hand side of Equation (12). Therefore the




FIG. 2: The correlation functions are given by contour integrals which circumscribe the four poles. The residue theorem states that the integral equals the sum over the four residues.

appropriate and only choice to define self-consistency equations is:

$$\langle S^y(0)_i S^y_i \rangle = \frac{1}{4} \quad (27)$$

In order to establish the link between the correlation functions and the corresponding Green's function the following spectral expansion of the Green's function at absolute zero as is readily obtained from Equation (3):

$$\begin{aligned} G_{ij\text{Ret}/\text{Adv}}^{yy}(\omega) &= \sum_n \frac{\langle n_0 | S_i^y | n \rangle \langle n | S_j^y | n_0 \rangle}{\omega - (\omega_n - \omega_{n_0}) \pm i\epsilon} \\ &\quad - \sum_n \frac{\langle n_0 | S_j^y | n \rangle \langle n | S_i^y | n_0 \rangle}{\omega + (\omega_n - \omega_{n_0}) \pm i\epsilon} \end{aligned} \quad (28)$$

where we have used that:

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(\omega - \omega')t} \theta(t) dt = \lim_{\epsilon \rightarrow +0} \frac{1}{2\pi} \frac{i}{\omega - \omega' + i\epsilon} \quad (29)$$

Here $|n_0\rangle$ is the ground state of the system and $|n\rangle$ refers to the complete set of eigenstates. The spectral representation (Equation (28)) shows that the integral:

$$\begin{aligned} &\int_{-\infty}^{\infty} d\omega e^{i\omega t} (G_{ij\text{Adv}}^{yy}(\omega) - G_{ij\text{Ret}}^{yy}(\omega)) \\ &= \lim_{\epsilon \rightarrow 0} \int_{-\infty}^{\infty} d\omega e^{i\omega t} (G_{ij}^{yy}(\omega + i\epsilon) - G_{ij}^{yy}(\omega - i\epsilon)) \end{aligned} \quad (30)$$

is a contour integral enclosing the poles of the Fourier transformed Green's function, as can be seen in Figure (2). The path in the upper and lower half planes correspond to the retarded and Green's functions respectively. According to the residue theorem of complex

analysis the value of the integral is given by the sum of the residues of the enclosed poles. As ω_{n_0} is by definition the lowest energy state, the spectral representation of the Green's function shows that the correlation function $\langle S_i^y(t) S_j^y \rangle$ corresponds to all negative poles and the conjugated correlations function $\langle S_j^y S_i^y(t) \rangle$ corresponds to all positive poles. Hence we obtain the desired correlation function $\langle S_i^y(t) S_j^y \rangle$ by restricting the contour integral to negative frequencies:

$$\begin{aligned} \langle S_i^y(t) S_j^y \rangle &= \\ &-i \lim_{\epsilon \rightarrow 0} \int_{-\infty}^0 e^{i\omega t} G_{ij}^{yy}(\omega + i\epsilon) - G_{ij}^{yy}(\omega - i\epsilon) d\omega \\ &+ \langle n_0 | S_i^y | n_0 \rangle \langle n_0 | S_j^y | n_0 \rangle \end{aligned} \quad (31)$$

The second term on the right side appears because its contribution in the spectral representation (Equation (28)) is canceled out. As we have broken the U(1) symmetry of the ground state in a way that $\langle S^y \rangle = 0$ this term will yield zero anyway and we will disregard it in further discussion. Using the Green's function's Fourier transform into k-space we obtain:

$$\begin{aligned} \langle S_i^y(t) S_j^y \rangle &= \\ &-i \lim_{\epsilon \rightarrow 0} \int \int_{-\infty}^0 d^3k d\omega e^{i(kr_{ij} + \omega t)} [G^{yy}(k, \omega + i\epsilon) \\ &- G^{yy}(k, \omega - i\epsilon)] d\omega \\ &= 2\pi \int d^3k e^{ikr_{ij}} [\text{Residue}(G^{yy}(k, \omega) e^{i\omega t}, -\omega_1) \\ &+ \text{Residue}(G^{yy}(k, \omega) e^{i\omega t}, -\omega_2)] \end{aligned} \quad (32)$$

Now we can define two self-consistency equation which determine the spin fields:

$$\begin{aligned} F_A &:= 2\pi \int d^3k [\text{Residue}(G_A^{yy}(k, \omega), -\omega_1) \\ &+ \text{Residue}(G_A^{yy}(k, \omega), -\omega_2)] - \frac{1}{4} = 0 \\ F_B &:= 2\pi \int d^3k [\text{Residue}(G_B^{yy}(k, \omega), -\omega_1) \\ &+ \text{Residue}(G_B^{yy}(k, \omega), -\omega_2)] - \frac{1}{4} = 0 \end{aligned} \quad (33)$$

In the canted anti-ferromagnetic and canted ferromagnetic phases we use Equation (19) to eliminate $\langle S_A^z \rangle$ and $\langle S_B^z \rangle$ and re-introduce h^z instead and obtain:

$$F_A^c(\langle S_A^x \rangle, \langle S_B^x \rangle, h^z) = 0$$

$$F_B^c(\langle S_A^x \rangle, \langle S_B^x \rangle, h^z) = 0 \quad (34)$$

similarly, we obtain the self-consistency equations for the ferromagnetic and anti-ferromagnetic phases:

$$\begin{aligned} F_A^{nc}(\langle S_A^z \rangle, \langle S_B^z \rangle, h^z) &= 0 \\ F_B^{nc}(\langle S_A^z \rangle, \langle S_B^z \rangle, h^z) &= 0 \end{aligned} \quad (35)$$

the three dimensional numerical integral contained in those equations can be reduced by one dimension by introducing a two dimensional density of states $\rho(\gamma'_1, \gamma'_2) = \int d^3k \delta(\gamma'_1 - \gamma_1(k)) \delta(\gamma'_2 - \gamma_2(k))$, so that:

$$\int d^3k \rightarrow \int d\gamma_1 d\gamma_2 \rho(\gamma_1, \gamma_2) \quad (36)$$

Consequently, in the self-consistency equations the DOS $\rho(\gamma_1, \gamma_2)$ is the only remaining term depending on the specific lattice structure. (In the actual numerical calculation it is, in order to avoid singularities in the origin, feasible to integrate over $\rho(\arccos(\gamma_1), \arccos(\gamma_2))$. For small k , $\arccos(\gamma_1)$ and $\arccos(\gamma_2)$ vary as $\sim k$ and therefore take the important contributions at small k more accurately into account than γ_1 and γ_2 which vary as $\sim k^2$.) Through this procedure we have archived a wider applicability of the corresponding equations as other systems, i.e. lattice geometries exhibiting frustration or two dimensional systems where linear spin-waves are still a valid approximation, can easily be accomplished by simply inserting the appropriate DOS. The DOS for the bipartite bcc lattice is shown in Figure 3.

IV. SELF-CONSISTENCY EQUATIONS AT FINITE TEMPERATURE

At finite temperature the system additionally exhibits thermal fluctuations expressed thorough averaging with the Boltzmann weight. As a consequence the negative and positive poles of the Green's functions do no longer refer solely to the corresponding correlations functions or their conjugates respectively. As the temperature is turned on the contribution of the poles to the correlation functions starts to overlap with the contribution of their conjugate correlation functions, where the degree of overlap is determined by the Boltzmann weight. Therefore the extraction of the correlations function from the corresponding commutator Green's function is not as straight forward as in the zero temperature case but the overlap to be reversed, i.e. an additional factor cancels out the contribution of the conjugate




FIG. 3: Two dimensional density of state $\rho(\gamma_1, \gamma_2)$ for the bipartite bcc lattice. The invariance of the DOS under $\gamma_1 \rightarrow -\gamma_1$ reflects the discrete translational symmetry of the bcc lattice along the diagonal, i.e. $\{x, y, z\} \rightarrow \{x + a/2, y + a/2, z + a/2\}$. The main contribution to the density of states comes from a relatively small area around three lines in the shape of the letter Y with a peak at $\gamma_1 = 0$ and $\gamma_2 = -0.327$.

function. This was done by Tyablikov [9, 10] in 1959. Here we follow a slightly different derivation.

As we have seen in the disquisition on the zero-temperature regime the adequate self-consistency equations are expressed by:

$$\langle S^y(0)_i S_i^y \rangle = \frac{1}{4} \quad (37)$$

The spectral representation of the Green's function at finite temperature is according to Equation (3) given by:

$$\begin{aligned} G_{ij\text{Ret/Adv}}^{yy}(\omega) = & \sum_{m,n} \frac{1}{2\pi} \frac{\langle n|S_i^y|m\rangle \langle m|S_j^y|n\rangle}{\omega - (\omega_m - \omega_n) \pm i\epsilon} \frac{e^{-\beta\omega_n}}{Z} \\ & - \sum_{m,n} \frac{1}{2\pi} \frac{\langle n|S_j^y|m\rangle \langle m|S_i^y|n\rangle}{\omega + (\omega_m - \omega_n) \pm i\epsilon} \frac{e^{-\beta\omega_n}}{Z} \end{aligned}$$

$$= \sum_{m,n} \frac{1}{2\pi} \frac{\langle n|S_i^y|m\rangle\langle m|S_j^y|n\rangle}{\omega - (\omega_m - \omega_n) \pm i\epsilon} \frac{e^{-\beta\omega_n} - e^{-\beta\omega_m}}{Z} \quad (38)$$

where we once again used that:

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(\omega-\omega')t} \theta(t) dt = \lim_{\epsilon \rightarrow +0} \frac{1}{2\pi} \frac{i}{\omega - \omega' + i\epsilon} \quad (39)$$

$|m\rangle$ and $|n\rangle$ are eigenstates of the Hamiltonian and Z refers to the partition function.

The Fourier transform of the correlation function is given by:

$$\begin{aligned} \langle S_i^y(t) S_j^y \rangle_{\omega} &= \sum_{m,n} \langle n|S_i^y|m\rangle\langle m|S_j^y|n\rangle \frac{e^{-\beta\omega_n}}{Z} \\ &\quad \times \delta(\omega - (\omega_m - \omega_n)) \end{aligned} \quad (40)$$

The relation to the Green's function is readily obtained by the following rearrangements:

$$\begin{aligned} \langle S_i^y(t) S_j^y \rangle_{\omega} &= \sum_{m,n'} \frac{\langle n|S_i^y|m\rangle\langle m|S_j^y|n\rangle}{Z} e^{-\beta\omega_n} \\ &\quad \times \frac{e^{\beta(\omega_n - \omega_m)} - 1}{e^{\beta(\omega_n - \omega_m)} - 1} \delta(\omega - (\omega_m - \omega_n)) \\ &\quad + \sum_n \frac{\langle n|S_i^y|n\rangle\langle n|S_j^y|n\rangle}{Z} e^{-\beta\omega_n} \\ &= \frac{\sum_{m,n'} \langle n|S_i^y|m\rangle\langle m|S_j^y|n\rangle}{e^{\beta\omega} - 1} \frac{e^{-\beta\omega_m} - e^{-\beta\omega_n}}{Z} \\ &\quad \times \delta(\omega - (\omega_m - \omega_n)) \\ &\quad + \sum_n \frac{\langle n|S_i^y|n\rangle\langle n|S_j^y|n\rangle}{Z} e^{-\beta\omega_n} \end{aligned} \quad (41)$$

The prime ' on the sum excludes terms where $m = n$ and was introduced to avoid singularities in the denominator. The omitted terms are accounted for in the second term of the RHS.

The following symbolic identity is valid for real ω

$$\lim_{\epsilon \rightarrow +0} \left[\frac{1}{\omega - \omega' + i\epsilon} - \frac{1}{\omega - \omega' - i\epsilon} \right] = -2\pi i \delta(\omega - \omega') \quad (42)$$

and immediately yields:

$$\begin{aligned} \langle S_i^y(t) S_j^y \rangle &= \\ &\quad i \lim_{\epsilon \rightarrow 0} \int_{-\infty}^0 e^{i\omega t} \frac{G_{ij}^{yy}(\omega + i\epsilon) - G_{ij}^{yy}(\omega - i\epsilon)}{e^{\beta\omega} - 1} d\omega \\ &\quad + \sum_n \frac{\langle n|S_i^y|n\rangle\langle n|S_j^y|n\rangle}{Z} e^{-\beta\omega_n} \end{aligned} \quad (43)$$

As pointed out by Stevens and Toombs [13] the second term on the right side is generally difficult to calculate. In principle it would have been possible to eliminate this term by also calculating the anti-commutator Green's function. Once again, as we have broken the U(1) symmetry of the ground state in a way such that $\langle S^y \rangle = 0$ this term will yield zero.

Using the Green's function's Fourier transform into k-space we obtain:

$$\begin{aligned}
\langle S_i^y(t) S_j^y \rangle &= \\
& i \lim_{\epsilon \rightarrow 0} \int \int_{-\infty}^0 d^3 k d\omega \frac{e^{i(kr_{ij} + \omega t)}}{e^{\beta\omega} - 1} \\
& [G^{yy}(k, \omega + i\epsilon) - G^{yy}(k, \omega - i\epsilon)] d\omega \\
& = -2\pi \int d^3 k e^{ikr_{ij}} \sum_{\omega_p = \omega_1, \omega_2} [\text{Residue} \left(\frac{G^{yy}(k, \omega) e^{i\omega t}}{e^{\beta\omega} - 1}, \omega_p \right) \\
& + \text{Residue} \left(\frac{G^{yy}(k, \omega) e^{i\omega t}}{e^{\beta\omega} - 1}, -\omega_p \right)]
\end{aligned} \tag{44}$$

Now we can define two self-consistency equations which determine the spin fields at finite temperature:

$$\begin{aligned}
F_A &:= \int d^3 k \sum_{\omega_p} [\text{Residue} \left(\frac{G_A^{yy}(k, \omega)}{e^{\beta\omega} - 1}, \omega_p \right) \\
& + \text{Residue} \left(\frac{G_A^{yy}(k, \omega)}{e^{\beta\omega} - 1}, -\omega_p \right)] - \frac{1}{4} = 0 \\
F_B &:= \int d^3 k \sum_{\omega_p} [\text{Residue} \left(\frac{G_B^{yy}(k, \omega)}{e^{\beta\omega} - 1}, \omega_p \right) \\
& + \text{Residue} \left(\frac{G_B^{yy}(k, \omega)}{e^{\beta\omega} - 1}, -\omega_p \right)] - \frac{1}{4} = 0
\end{aligned} \tag{45}$$

Using the DOS of Equation (36) those are written as:

$$\begin{aligned}
F_A &= \int d\gamma_1 d\gamma_2 \rho(\gamma_1, \gamma_2) \times \\
& \sum_{\omega_p} [\text{Residue} \left(\frac{G_A^{yy}(\gamma_1, \gamma_2, \omega)}{e^{\beta\omega} - 1}, \omega_p \right) \\
& + \text{Residue} \left(\frac{G_A^{yy}(\gamma_1, \gamma_2, \omega)}{e^{\beta\omega} - 1}, -\omega_p \right)] - \frac{1}{4} = 0 \\
F_B &= \int d\gamma_1 d\gamma_2 \rho(\gamma_1, \gamma_2) \times \\
& \sum_{\omega_p} [\text{Residue} \left(\frac{G_B^{yy}(\gamma_1, \gamma_2, \omega)}{e^{\beta\omega} - 1}, \omega_p \right) \\
& + \text{Residue} \left(\frac{G_B^{yy}(\gamma_1, \gamma_2, \omega)}{e^{\beta\omega} - 1}, -\omega_p \right)] - \frac{1}{4} = 0
\end{aligned} \tag{46}$$

Again, in the canted anti-ferromagnetic and canted ferromagnetic phases we use the mean-field equations [2, 3, 4] to eliminate $\langle S_A^z \rangle$ and $\langle S_B^z \rangle$ and obtain:

$$\begin{aligned} F_A^c(\langle S_A^x \rangle, \langle S_B^x \rangle, h^z, T) &= 0 \\ F_B^c(\langle S_A^x \rangle, \langle S_B^x \rangle, h^z, T) &= 0 \end{aligned} \quad (47)$$

Similarly, we obtain the self-consistency equations for the ferromagnetic and anti-ferromagnetic phases:

$$\begin{aligned} F_A^{nc}(\langle S_A^z \rangle, \langle S_B^z \rangle, h^z, T) &= 0 \\ F_B^{nc}(\langle S_A^z \rangle, \langle S_B^z \rangle, h^z, T) &= 0 \end{aligned} \quad (48)$$

V. GREEN'S FUNCTIONS IN EXPLICIT FORM

In the remaining three Chapters we give the Green's functions and the corresponding correlations functions and carry out the derivation of the matrix equation defining the Green's functions.

We start with the Green's functions for the anisotropic Heisenberg model for the canted phases, i.e. for the canted ferromagnetic and the canted anti-ferromagnetic phases

$$\begin{aligned} G_A^{xy}(k, \omega) &= \frac{-i\omega \langle S_A^z \rangle}{(2\pi)^4(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2)} \times \\ &(\omega^2 + 4CD + 4DA_1 - 4(C + A_2)(B_2 + A_2(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1))) \\ G_B^{xy}(k, \omega) &= \frac{-i\omega \langle S_B^z \rangle}{(2\pi)^4(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2)} \times \\ &(\omega^2 + 4CD + 4DA_2 - 4(C + A_1)(B_1 + A_1(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1))) \end{aligned} \quad (49)$$

$$\begin{aligned} G_A^{yy}(k, \omega) &= \frac{-1}{(2\pi)^4(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2)} \times \\ &(D(\omega^2 + 4CD + 4DA_2) + (B_1 + A_1(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1)) \\ &(\omega^2 - 4(C + A_2)(B_2 + A_2(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1)))) \\ G_B^{yy}(k, \omega) &= \frac{-1}{(2\pi)^4(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2)} \times \\ &(D(\omega^2 + 4CD + 4DA_1) + (B_1 + A_1(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1)) \\ &(\omega^2 - 4(C + A_1)(B_1 + A_1(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1)))) \end{aligned} \quad (50)$$

$$\begin{aligned}
G_A^{zy}(k, \omega) &= \frac{-i\omega \langle S_A^x \rangle}{(2\pi)^4(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2)} \times \\
&\quad (\omega^2 + 4CD + 4DA_1 - 4(C + A_2)(B_2 + A_2(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1))) \\
G_B^{zy}(k, \omega) &= \frac{-i\omega \langle S_B^x \rangle}{(2\pi)^4(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2)} \times \\
&\quad (\omega^2 + 4CD + 4DA_2 - 4(C + A_1)(B_1 + A_1(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1)))
\end{aligned} \tag{51}$$

The poles are given by:

$$\begin{aligned}
\omega_{1/2} &= 2(A_1(B_1 + A_1(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1)) + \\
&\quad A_2(B_2 + A_2(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1))) - 4CD \\
&\quad \pm 2[(-2CD + A_1(B_1 + A_1(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1)) \\
&\quad + A_2(B_2 + A_2(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1)))^2 \\
&\quad - 4(C^2 - A_1A_2)(D^2 - (B_1 + A_1(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1)) \\
&\quad \times (B_2 + A_2(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1)))]^{1/2}
\end{aligned} \tag{52}$$

All parameter A_1, A_2, B_1, B_2, C and D are as stated in Equation (21).

As condition (19) does not apply for the ferromagnetic nor for the anti-ferromagnetic states, these Green's functions are structurally different:

$$\begin{aligned}
G_A^{xy}(k, \omega) &= \frac{i\omega((A_3 + A_4)B_3\langle S_B^z \rangle - (-^2\omega^2 + A_4^2 + B_3B_4)\langle S_A^z \rangle)}{(2\pi)^4(^2\omega^2 - ^2\omega_1^2)(^2\omega^2 - ^2\omega_2^2)} \\
G_B^{xy}(k, \omega) &= \frac{i\omega((A_3 + A_4)B_4\langle S_A^z \rangle - (-^2\omega^2 + A_3^2 + B_3B_4)\langle S_B^z \rangle)}{(2\pi)^4(^2\omega^2 - ^2\omega_1^2)(^2\omega^2 - ^2\omega_2^2)}
\end{aligned} \tag{53}$$

$$\begin{aligned}
G_A^{yy}(k, \omega) &= \frac{1}{(2\pi)^4(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2)} \times \\
&\quad ((\omega^2 A_3 - A_3A_4^2 + A_4B_3B_4)\langle S_A^z \rangle + B_3(\omega^2 + A_3A_4 - B_3B_4)\langle S_B^z \rangle) \\
G_B^{yy}(k, \omega) &= \frac{1}{(2\pi)^4(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2)} \times \\
&\quad (B_4(^2\omega^2 + A_3A_4 - B_3B_4)\langle S_A^z \rangle + (^2\omega^2 A_4 - A_3^2A_4 + A_3B_3B_4)\langle S_B^z \rangle)
\end{aligned} \tag{54}$$

with

$$\omega_{1/2} = \frac{A_3 + A_4 \pm \sqrt{(A_3 - A_4)^2 + 4B_3B_4}}{2} \tag{55}$$

$$\begin{aligned}
A_3 &= 2h_z + 4\langle S_A^z \rangle (J_2^\parallel - J_2^\top \gamma_2(k)) + 4\langle S_B^z \rangle J_1^\parallel \\
A_4 &= 2h_z + 4\langle S_B^z \rangle (J_2^\parallel - J_2^\top \gamma_2(k)) + 4\langle S_A^z \rangle J_1^\parallel \\
B_3 &= -4\langle S_A^z \rangle J_1^\top \gamma_1(k) \\
B_4 &= -4\langle S_B^z \rangle J_1^\top \gamma_1(k)
\end{aligned} \tag{56}$$

A. The Correlation Functions in Explicit Form

Here we state the relevant correlation functions. All other correlation functions yield identically $\frac{1}{4}\langle S_{x_{A/B}} \rangle$ or $\frac{1}{4}\langle S_{z_{A/B}} \rangle$ and are therefore not suitable as self-consistency equations. We start with the Green's function for the canted anti-ferromagnetic and the canted ferromagnetic phase at finite temperature:

$$\begin{aligned}
\langle S_i^y S_{j \in A}^y(0) \rangle_k &= -\frac{\coth\left(\frac{\beta\omega_1}{2}\right)}{2(\omega_1^3 - \omega_1\omega_2^2)} \times \\
&\left(4CD^2 + \omega_1^2 D + \omega_1^2 B_1 - 4CB_1 B_2 \right. \\
&- 4A_2^2 B_1 (4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1) \\
&+ A_1 (4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1) \times \\
&(\omega_1^2 - 4CB_2 - 4A_2^2 (4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1) \\
&- 4A_2 (B_2 + C(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1))) \\
&\left. + 4A_2 (D^2 - B_1 (B_2 + C(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1))) \right) \\
&- \frac{\coth\left(\frac{\beta\omega_2}{2}\right)}{2(\omega_2^3 - \omega_1^2\omega_2)} \left(4CD^2 + \omega_2^2 D + \omega_2^2 B_1 - 4CB_1 B_2 \right. \\
&- 4A_2^2 B_1 (4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1) \\
&+ A_1 (4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1) \times \\
&(\omega_2^2 - 4CB_2 - 4A_2^2 (4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1) \\
&- 4A_2 (B_2 + C(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1))) \\
&\left. + 4A_2 (D^2 - B_1 (B_2 + C(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1))) \right)
\end{aligned} \tag{57}$$

$$\begin{aligned}
\langle S_i^y S_{j \in B}^y(0) \rangle_k &= -\frac{\coth\left(\frac{\beta\omega_1}{2}\right)}{2(\omega_1^3 - \omega_1\omega_2^2)} \times \\
&\left(4CD^2 + \omega_1^2 D + \omega_1^2 B_2 - 4CB_2 B_1 \right. \\
&- 4A_1^2 B_2 (4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1)
\end{aligned}$$

$$\begin{aligned}
& +A_2(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1) \times \\
& (\omega_1^2 - 4CB_1 - 4A_1^2(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1) \\
& - 4A_1(B_1 + C(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1))) + \\
& 4A_1(D^2 - B_2(B_1 + C(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1)))) \\
& - \frac{\coth\left(\frac{\beta\omega_2}{2}\right)}{2(\omega_2^3 - \omega_1^2\omega_2)} \left(4CD^2 + \omega_2^2D + \omega_2^2B_2 - 4CB_2B_1 \right. \\
& - 4A_2^2B_2(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1) \\
& + A_1(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1) \times \\
& (\omega_2^2 - 4CB_1 - 4A_1^2(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1) \\
& - 4A_1(B_1 + C(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1))) \\
& \left. + 4A_1(D^2 - B_2(B_1 + C(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1)))) \right) \tag{58}
\end{aligned}$$

At zero temperature these functions become:

$$\begin{aligned}
\langle S_i^y S_{j \in A}^y(0) \rangle_k &= -\frac{1}{2\sqrt{\omega_1^2}(\omega_1^2 - \omega_2^2)} \times \\
& \left(4CD^2 + \omega_1^2D + \omega_1^2B_1 - 4CB_1B_2 \right. \\
& - 4A_2^2B_1(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1) \\
& + A_1(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1) \times \\
& (\omega_1^2 - 4CB_2 - 4A_2^2(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1) \\
& - 4A_2(B_2 + C(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1))) \\
& \left. + 4A_2(D^2 - B_1(B_2 + C(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1)))) \right) \\
& - \frac{1}{2\sqrt{\omega_2^2}(\omega_2^2 - \omega_1^2)} \left(4CD^2 + \omega_2^2D + \omega_2^2B_1 - 4CB_1B_2 \right. \\
& - 4A_2^2B_1(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1) \\
& + A_1(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1) \times \\
& (\omega_2^2 - 4CB_2 - 4A_2^2(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1) \\
& - 4A_2(B_2 + C(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1))) \\
& \left. + 4A_2(D^2 - B_1(B_2 + C(4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1)))) \right) \tag{59}
\end{aligned}$$

$$\langle S_i^y S_{j \in B}^y(0) \rangle_k = -\frac{1}{2\sqrt{\omega_1^2}(\omega_1^2 - \omega_2^2)} \times$$

$$\begin{aligned}
& \left(4CD^2 + \omega_1^2 D + \omega_1^2 B_2 - 4CB_2B_1 \right. \\
& - 4A_1^2 B_2 (4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1) \\
& + A_2 (4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1) \times \\
& (\omega_1^2 - 4CB_1 - 4A_1^2 (4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1)) \\
& - 4A_1 (B_1 + C(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1))) + \\
& \left. 4A_1 (D^2 - B_2 (B_1 + C(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1))) \right) \\
& - \frac{1}{2\sqrt{\omega_2^2}(\omega_2^2 - \omega_1^2)} \left(4CD^2 + \omega_2^2 D + \omega_2^2 B_2 - 4CB_2B_1 \right. \\
& - 4A_2^2 B_2 (4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1) \\
& + A_1 (4\langle S_B^x \rangle^2 + 4\langle S_B^z \rangle^2 - 1) \times \\
& (\omega_2^2 - 4CB_1 - 4A_1^2 (4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1)) \\
& - 4A_1 (B_1 + C(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1))) \\
& \left. + 4A_1 (D^2 - B_2 (B_1 + C(4\langle S_A^x \rangle^2 + 4\langle S_A^z \rangle^2 - 1))) \right) \tag{60}
\end{aligned}$$

the correlation functions for the ferromagnetic and the anti-ferromagnetic phases are:

$$\begin{aligned}
\langle S_i^y S_{j \in A}^y(0) \rangle_k &= \\
& \frac{\coth(\frac{\beta\omega_1}{2})}{2(\omega_1^3 - \omega_1\omega_2^2)} \times \\
& ((A_3\langle S_A^z \rangle + B_3\langle S_B^z \rangle)\omega_1^2 - (A_3A_3 - B_3B_4)(A_4\langle S_A^z \rangle - B_3\langle S_B^z \rangle)) \\
& + \frac{\coth(\frac{\beta\omega_2}{2})}{2(\omega_2^3 - \omega_1^2\omega_2)} \times \\
& ((A_3\langle S_A^z \rangle + B_3\langle S_B^z \rangle)\omega_2^2 - (A_3A_4 - B_3B_4)(A_4\langle S_A^z \rangle - B_3\langle S_B^z \rangle)) \tag{61}
\end{aligned}$$

$$\begin{aligned}
\langle S_i^y S_{j \in B}^y(0) \rangle_k &= \\
& \frac{\coth(\frac{\beta\omega_1}{2})}{2(\omega_1^3 - \omega_1\omega_2^2)^4} \times \\
& ((A_4\langle S_B^z \rangle + B_4\langle S_A^z \rangle)\omega_1^2 - (A_4A_4 - B_4B_3)(A_3\langle S_B^z \rangle - B_4\langle S_A^z \rangle)) \\
& + \frac{\coth(\frac{\beta\omega_2}{2})}{2(\omega_2^3 - \omega_1^2\omega_2)} \times \\
& ((A_4\langle S_B^z \rangle + B_4\langle S_A^z \rangle)\omega_2^2 - (A_4A_3 - B_4B_3)(A_3\langle S_B^z \rangle - B_4\langle S_A^z \rangle)) \tag{62}
\end{aligned}$$

which can be written in the zero temperature limit as:

$$\langle S_i^y S_{j \in A}^y(0) \rangle_k =$$

$$\frac{(A_3A_4 - B_3B_4)(A_4\langle S_{z_A}\rangle - B_3\langle S_{z_B}\rangle) + (A_3\langle S_{z_A}\rangle - B_3\langle S_{z_B}\rangle)\sqrt{(\omega_1\omega_2)^2}}{\sqrt{(\omega_1\omega_2)^2}(\sqrt{\omega_1^2} + \sqrt{\omega_2^2})} \quad (63)$$

$$\begin{aligned} \langle S_i^y S_{j \in B}^y(0) \rangle_k = \\ \frac{(A_3A_4 - B_3B_4)(A_3\langle S_{z_B}\rangle - B_4\langle S_{z_A}\rangle) + (A_4\langle S_{z_B}\rangle - B_4\langle S_{z_A}\rangle)\sqrt{(\omega_1\omega_2)^2}}{\sqrt{(\omega_1\omega_2)^2}(\sqrt{\omega_1^2} + \sqrt{\omega_2^2})} \end{aligned} \quad (64)$$

B. Derivation of the Matrix Equation

Finally we carry out the derivation of the matrix equation, evolving from the equations of motion. The three equations of motion for the Green's functions after employing the cumulant decoupling scheme become:

$$\begin{aligned} i\partial_t G_{ij}^{xy}(t) &= i\delta(t)\delta_{ij}\langle S_i^z(t) \rangle - ih_z G_{ij}^{yy}(t) \\ &\quad - 2i \sum_l J_{il}^{\parallel} (\langle S_i^y(t) \rangle G_{lj}^{zy}(t) + \langle S_l^z(t) \rangle G_{ij}^{yy}(t)) \\ &\quad + 2i \sum_l J_{il}^{\top} (\langle S_i^z(t) \rangle G_{lj}^{yy}(t) + \langle S_l^y(t) \rangle G_{ij}^{zy}(t)) \end{aligned} \quad (65)$$

$$\begin{aligned} i\partial_t G_{ij}^{yy}(t) &= ih_z G_{ij}^{xy}(t) \\ &\quad + 2i \sum_l J_{il}^{\parallel} (\langle S_i^x(t) \rangle G_{lj}^{zy}(t) + \langle S_l^z(t) \rangle G_{ij}^{xy}(t)) \\ &\quad - 2i \sum_l J_{il}^{\top} (\langle S_i^z(t) \rangle G_{lj}^{xy}(t) + \langle S_l^x(t) \rangle G_{ij}^{zy}(t)) \end{aligned} \quad (66)$$

$$\begin{aligned} i\partial_t G_{ij}^{zy}(t) &= -i\delta(t)\delta_{ij}\langle S_i^x(t) \rangle \\ &\quad + 2i \sum_l J_{il}^{\top} (\langle S_i^y(t) \rangle G_{lj}^{xy}(t) + \langle S_l^x(t) \rangle G_{ij}^{yy}(t)) \\ &\quad - 2i \sum_l J_{il}^{\parallel} (\langle S_i^x(t) \rangle G_{lj}^{yy}(t) + \langle S_l^y(t) \rangle G_{ij}^{xy}(t)) \end{aligned} \quad (67)$$

Fourier transforming into ω -space and assume that the spins are constants in time:

$$\begin{aligned} \omega G_{ij}^{xy}(\omega) &= \frac{i}{2\pi} \delta_{ij} \langle S_i^z \rangle - ih_z G_{ij}^{yy}(\omega) \\ &\quad - 2i \sum_l J_{il}^{\parallel} (\langle S_i^y \rangle G_{lj}^{zy}(\omega) + \langle S_l^z \rangle G_{ij}^{yy}(\omega)) \\ &\quad + 2i \sum_l J_{il}^{\top} (\langle S_i^z \rangle G_{lj}^{yy}(\omega) + \langle S_l^y \rangle G_{ij}^{zy}(\omega)) \end{aligned} \quad (68)$$

$$\begin{aligned}
\omega G_{ij}^{yy}(\omega) &= ih_z G_{ij}^{xy}(\omega) \\
&+ 2i \sum_l J_{il}^{\parallel} (\langle S_i^x \rangle G_{lj}^{zy}(\omega) + \langle S_l^z \rangle G_{ij}^{xy}(\omega)) \\
&- 2i \sum_l J_{il}^{\top} (\langle S_i^z \rangle G_{lj}^{xy}(\omega) + \langle S_l^x \rangle G_{ij}^{zy}(\omega))
\end{aligned} \tag{69}$$

$$\begin{aligned}
\omega G_{ij}^{zy}(\omega) &= -\frac{i}{2\pi} \delta_{ij} \langle S_i^x \rangle \\
&+ 2i \sum_l J_{il}^{\top} (\langle S_i^y \rangle G_{lj}^{xy}(\omega) + \langle S_l^x \rangle G_{ij}^{yy}(\omega)) \\
&- 2i \sum_l J_{il}^{\parallel} (\langle S_i^x \rangle G_{lj}^{yy}(\omega) + \langle S_l^y \rangle G_{ij}^{xy}(\omega))
\end{aligned} \tag{70}$$

In the canted anti-ferromagnetic and the anti-ferromagnetic phases the spins on the two sub-lattices assume different values. Therefore we need to split up every Green's function before we can Fourier Transform into k-space. We define $G_A^{\mu\nu}(k, \omega)$ and $G_B^{\mu\nu}(k, \omega)$. $G_A^{\mu\nu}(k, \omega)$ refers to $G_{ij}^{\mu\nu}(\omega)$ when the site i is on sub-lattice A and $G_B^{\mu\nu}(k, \omega)$ when i is on sub-lattice B.

$$\begin{aligned}
\omega G_{A_k}^{xy}(\omega) &= \frac{i}{(2\pi)^4} \langle S_A^z \rangle - 2ih_z G_{A_k}^{yy}(\omega) \\
&- 4i \langle S_A^y \rangle [G_{A_k}^{zy}(\omega) J_2^{\parallel} \gamma_2(k) + G_{B_k}^{zy}(\omega) J_1^{\parallel} \gamma_1(k)] \\
&- 4i G_{A_k}^{yy}(\omega) [\langle S_A^z \rangle J_2^{\parallel} \gamma_2(0) + \langle S_B^z \rangle J_1^{\parallel} \gamma_1(0)] \\
&+ 4i \langle S_A^z \rangle [G_{A_k}^{yy}(\omega) J_2^{\top} \gamma_2(k) + G_{B_k}^{yy}(\omega) J_1^{\top} \gamma_1(k)] \\
&+ 4i G_{A_k}^{zy}(\omega) [\langle S_A^y \rangle J_2^{\top} \gamma_2(0) + \langle S_B^y \rangle J_1^{\top} \gamma_1(0)]
\end{aligned} \tag{71}$$

$$\begin{aligned}
\omega G_{A_k}^{yy}(\omega) &= -2ih_z G_{A_k}^{xy}(\omega) \\
&+ 4i \langle S_A^x \rangle [G_{A_k}^{zy}(\omega) J_2^{\parallel} \gamma_2(k) + G_{B_k}^{zy}(\omega) J_1^{\parallel} \gamma_1(k)] \\
&+ 4i G_{A_k}^{xy}(\omega) [\langle S_A^z \rangle J_2^{\parallel} \gamma_2(0) + \langle S_B^z \rangle J_1^{\parallel} \gamma_1(0)] \\
&- 4i \langle S_A^z \rangle [G_{A_k}^{xy}(\omega) J_2^{\top} \gamma_2(k) + G_{B_k}^{xy}(\omega) J_1^{\top} \gamma_1(k)] \\
&- 4i G_{A_k}^{zy}(\omega) [\langle S_A^x \rangle J_2^{\top} \gamma_2(0) + \langle S_B^x \rangle J_1^{\top} \gamma_1(0)]
\end{aligned} \tag{72}$$

$$\begin{aligned}
\omega G_{A_k}^{zy}(\omega) &= -\frac{i}{(2\pi)^4} \langle S_A^x \rangle \\
&+ 4i \langle S_A^y \rangle [G_{A_k}^{xy}(\omega) J_2^{\top} \gamma_2(k) + G_{B_k}^{xy}(\omega) J_1^{\top} \gamma_1(k)]
\end{aligned}$$

$$\begin{aligned}
& -4iG_{A_k}^{yy}(\omega)[\langle S_A^x \rangle J_2^\top \gamma_2(0) + \langle S_B^x \rangle J_1^\top \gamma_1(0)] \\
& -4i\langle S_A^x \rangle [G_{A_k}^{yy}(\omega) J_2^\top \gamma_2(k) + G_{B_k}^{yy}(\omega) J_1^\top \gamma_1(k)] \\
& -4iG_{A_k}^{xy}(\omega)[\langle S_A^y \rangle J_2^\top \gamma_2(0) + \langle S_B^y \rangle J_1^\top \gamma_1(0)]
\end{aligned} \tag{73}$$

VI. CONCLUSIONS

In conclusion, we analysed the anisotropic Heisenberg model in an external field on the three dimensional bcc lattice by employing the well-established technique of real-time Green's functions for spin systems. The series of infinite order Green's functions as it appears in the equation of motion was truncated by applying cumulant decoupling and the resulting random phase approximation accounts for linear spin-waves. We are the first to apply this method to the canted anti-ferromagnetic phase entailing a set of six algebraic equations. The innate self-consistency equations herein constitute a three dimensional numerical integral over the k-space. By introducing a two dimensional density of states the integral was reduced to two dimensions where the lattice generating functions serve as new integration variables. In the appearing integrals the DOS is the only quantity that depends on the structure of the lattice. Hence, once the DOS is computed for a certain lattice geometry the further calculation remains unaltered. Therefore our method is widely applicable and easily adjustable to various magnetic systems where canted phases are in the center of interest. This also holds for two dimensional lattices where linear spin waves are expected to yield a reasonable approximation.

VII. ACKNOWLEDGEMENTS

One of the authors (Miklós Gulácsi) wishes to thank James L. Smith for his friendship, guidance and encouragement over the years. Thanks for the "*shmokos*"!

APPENDIX A: CONNECTION TO ⁴HE PHYSICS

For the interested reader, we briefly summarize the connection between the anisotropic Heisenberg model and ⁴He as introduced by Matsubara and Matsuda [14, 15], and used by Matsuda and Tsuneto [16], Fisher [5], Liu and Fisher [6] and most recently by Stoffel and

Gulácsi [2, 3, 4].

Apart from possible ^3He impurities ^4He is a bosonic system and the generic Hamiltonian for such systems in the language of second quantization is given by:

$$H = \int d^3x \hat{\psi}^\dagger(\mathbf{x}) \left(-\frac{1}{2m} \nabla^2 + \mu \right) \hat{\psi}(\mathbf{x}) + \frac{1}{2} \int d^3x d^3x' \hat{\psi}^\dagger(\mathbf{x}) \hat{\psi}^\dagger(\mathbf{x}') V(\mathbf{x} - \mathbf{x}') \hat{\psi}(\mathbf{x}) \hat{\psi}(\mathbf{x}') \quad (\text{A1})$$

where $\hat{\psi}^\dagger(\mathbf{x})$, the particle creation operator and $\hat{\psi}(\mathbf{x})$, the corresponding destruction operator obey the usual bosonic commutator relations. Hamiltonians in three dimensions such as in Equation (A1) are not solvable even for elementary potentials $V(\mathbf{x})$ such as the Dirac delta distribution. Therefore, further approximations has to be implemented. An approximation which proved particularly successful for the description of liquid Helium is know as the *quantum lattice gas* model and was first introduced by Matsubara and Matsuda [14, 15].

In the quantum lattice gas model one works with a space lattice of discrete lattice points rather than the continuum. This approximation proves to be very useful to study solid states as the spacial discretization of this model serves as a natural frame for the crystal lattice. Also in this model no specific knowledge of the density distribution of the atoms is needed.

According to Matsubara and Tsuneto [16] the generic Hamiltonian Equation (A1) in the discrete lattice model reads:

$$H = \mu \sum_i n_i + \sum_{ij} u_{ij} (a_i^\dagger - a_j^\dagger) (a_i - a_j) + \sum_{ij} V_{ij} n_i n_j \quad (\text{A2})$$

Here u_{ij} are non-zero for nearest neighbor and next nearest neighbor hopping and otherwise zero. The values of u_{nn} and u_{nnn} are such that the kinetic energy is isotropic up to the 4th order. In the case of a bcc lattice (two interpenetrating sc lattices) the matrix elements are given by:

$$u_{nn} = \frac{2}{3} \frac{1}{4ma^2} \quad (\text{A3})$$

$$u_{nnn} = \frac{1}{3} \frac{1}{4ma^2} \quad (\text{A4})$$

As the atoms do not penetrate each other there can exist only one atom at a time on a lattice site. Consequently $a^\dagger$ and a are the creation and annihilation operators of a hard

core boson commuting on different lattice sites:

$$[a_i^\dagger, a_j^\dagger]_- = [a_i, a_j]_- = [a_i, a_j^\dagger]_- = 0 \quad (i \neq j) \quad (\text{A5})$$

but obey the anti-commutator relations on identical sites:

$$\begin{aligned} [a_i^\dagger, a_i^\dagger]_+ &= [a_i, a_i]_+ = 0 \\ \left[a_i, a_i^\dagger \right]_+ &= 1 \end{aligned} \quad (\text{A6})$$

Equation (A2) is the Bose-Hubbard model in three dimensions for hard core bosons. Due to the unusual statistics of hard core bosons, Wick theorem cannot be applied and hence, the common formalism of perturbative field theory is not applicable. The way out is to transform the model to an equivalent spin model [14, 15], namely by using

$$\begin{aligned} a_j^\dagger &= S_j^x - iS_j^y \\ a_j &= S_j^x + iS_j^y \\ n_j &= \frac{1}{2} - S_j^z \end{aligned} \quad (\text{A7})$$

It can be verified that the usual Lie algebra for spin 1/2 particles preserves the mixed commutation/anti-commutation relations for hard-core bosons. This substitution transforms the hard-core bosonic Hubbard model into a spin model:

$$\begin{aligned} H &= \mu \sum_i \left(\frac{1}{2} - S_i^z \right) \\ &+ \sum_{ij} u_{ij} (1 - S_i^z - S_j^z - 2S_i^x S_j^x - 2S_i^y S_j^y) \\ &+ \sum_{ij} V_{ij} \left(\frac{1}{4} - \frac{S_i^z}{2} - \frac{S_j^z}{2} + S_i^z S_j^z \right) \end{aligned} \quad (\text{A8})$$

If we adjust the notation to conform with the usual standards of spin models, we re-obtain the anisotropic Heisenberg from Equation (1):

$$H = h^z \sum_i S_i^z + \sum_{ij} J_{ij}^\parallel S_i^z S_j^z + \sum_{ij} J_{ij}^\top (S_i^x S_j^x + S_i^y S_j^y) \quad (\text{A9})$$

with:

$$\begin{aligned} J_{ij}^\parallel &= V_{ij} \\ J_{ij}^\top &= -2u_{ij} \\ h^z &= -\mu + \sum_j J_{ij}^\top - \sum_j J_{ij}^\parallel \end{aligned} \quad (\text{A10})$$

If the above presented transformation is used for ^4He , then the values of the J 's also have to be chosen such as to mimic ^4He . The interactions between the ^4He atoms are controlled by van-der-Waals forces and their repulsive nature at very short distances determines negative nearest neighbor interaction $J_1^\parallel$, evoking anti-ferromagnetic ordering in the spin language. The corresponding Lennard-Jones potential is short ranged and therefore it is sufficient [6] to only consider nearest and next nearest neighbor interactions. Hence, for ^4He the J values will be $J_1^\parallel = -q_1 J_{i \in A, j \in B}^\parallel$, $J_2^\parallel = -q_2 J_{i \in A, j \in A}^\parallel$, $J_1^\top = -q_1 J_{i \in A, j \in B}^\top$ and $J_2^\top = -q_2 J_{i \in A, j \in A}^\top$, where $q_1 = 6$ and $q_2 = 8$ are the number of nearest and next nearest neighbors on the bipartite bcc lattice. Liu and Fisher [6] in their calculations used, $J_1^\top = 1.4K$, $J_2^\top = 0.5K$, $J_1^\parallel = -3.8K$ and $J_2^\parallel = -1.7K$. In a ^4He calculation the results do not change [3, 4] if the $J^\parallel$'s are within ± 2 range of these values and $J_B^\parallel > J_A^\parallel$ and $J^\top$'s values remain positive.

Defining two sub-lattices gave [5, 6, 14, 15, 16] a possibility to establish the diagonal long-range order of solids in a natural way: sub-lattice A represents the centers of the ^4He ions, hence it coincides with the ion lattice. Sub-lattice B defines the interstitials, the space in-between those atomic centers. In the liquid phases, of course, the occupation number on both sub-lattices is equal as there is no spacial density variation. In Table (I) we gave the various magnetic phases of the anisotropic Heisenberg model. These phases, however identify the corresponding phases of the ^4He system, as presented in Table (II).

APPENDIX B: THE MEAN-FIELD LIMIT

In the mean-field solution of Liu and Fisher [6] the Green's functions have not been explicitly evaluated, for the sake of completeness we re-derive these hereafter.

First we will re-derive the classical mean-field approximation as was pioneered by Liu and Fisher [6] and briefly state some key properties. We will further show that this approximation is a special case of the random-phase approximation.

The anisotropic Heisenberg Hamiltonian in the classical mean-field approximation is obtained by substituting the spin 1/2 operators with their respective expectation values:

$$\begin{aligned} H_{MF} = & -h^z (\langle S_A^z \rangle + \langle S_B^z \rangle) \\ & -2J_1^\parallel \langle S_A^z \rangle \langle S_B^z \rangle - J_2^\parallel (\langle S_A^z \rangle \langle S_A^z \rangle + \langle S_B^z \rangle \langle S_B^z \rangle) \\ & -2J_1^\top \langle S_A^x \rangle \langle S_B^x \rangle - J_2^\top (\langle S_A^x \rangle \langle S_A^x \rangle + \langle S_B^x \rangle \langle S_B^x \rangle) \end{aligned}$$

Spin Configuration	Magnetic Phase	ODLRO	DLRO	^4He -Phase
$\uparrow\uparrow$	FE	No	No	Normal Liquid
$\nearrow\nearrow$	CFE	Yes	No	Superfluid
$\nearrow\nwarrow$	CAF	Yes	Yes	Supersolid
$\uparrow\downarrow$	AF	No	Yes	Normal Solid

TABLE II: The phases of ^4He corresponding to the phases of the anisotropic Heisenberg model. Similarly to Table (I) the phases are defined by their long range order, i.e., off-diagonal long-range order (ODLRO) and diagonal long-range order (DLRO).

(B1)

Here $J_1^\parallel = -q_1 J_{i \in A, j \in B}^\parallel$, $J_2^\parallel = -q_2 J_{i \in A, j \in A}^\parallel$, $J_1^\top = -q_1 J_{i \in A, j \in B}^\top$ and $J_2^\top = -q_1 J_{i \in A, j \in A}^\top$ where $q_1 = 6$ and $q_2 = 8$ are the number of nearest and next nearest neighbours on the bipartite bcc lattice. The mean value of S_y drops out as the randomly broken symmetry $S_x \leftrightarrow S_y$ (off-diagonal long-range order) allows for $\langle S_y \rangle = 0$. The standard method of deriving the corresponding self-consistency equations is to minimize the Helmholtz's Free energy $F = H - TS$. The entropy S is given by the pseudo spin entropy of the system:

$$S = -\frac{1}{2} \left[\left(\frac{1}{2} + S_A \right) \ln \left(\frac{1}{2} + S_A \right) + \left(\frac{1}{2} - S_A \right) \ln \left(\frac{1}{2} - S_A \right) \right. \\ \left. + \left(\frac{1}{2} + S_B \right) \ln \left(\frac{1}{2} + S_B \right) + \left(\frac{1}{2} - S_B \right) \ln \left(\frac{1}{2} - S_B \right) \right] \quad (\text{B2})$$

where $S_A = \sqrt{\langle S_{z_A} \rangle^2 + \langle S_{x_A} \rangle^2}$ and $S_B = \sqrt{\langle S_{z_B} \rangle^2 + \langle S_{x_B} \rangle^2}$. We could equally well say that the state of the system is determined by minimizing the internal energy $\langle H \rangle$, subject to an additional constraint given by Equation (B2). In this picture the temperature becomes a Lagrange multiplier and at absolute zero, where $T = S = 0$ we obtain, as $\lim_{S_A, S_B \rightarrow 0} S = 0$,

$$\sqrt{\langle S_A^x \rangle^2 + \langle S_A^z \rangle^2} = \sqrt{\langle S_B^x \rangle^2 + \langle S_B^z \rangle^2} = \frac{1}{2} \quad (\text{B3})$$

This result implies that this approximation does not take quantum fluctuations into account.

In the canted anti-ferromagnetic and the canted ferromagnetic states there are four self-consistency equations in the ferromagnetic and anti-ferromagnetic phases; where $\langle S_x \rangle = \langle S_y \rangle = 0$ they are reduced in number by two. These equations are readily obtained by differentiating the free energy with respect to $\langle S_z \rangle$ and $\langle S_x \rangle$ respectively. The resulting equations can be rearranged to yield:

$$\begin{aligned}
\langle S_A^x \rangle &= \frac{2J_1^\top \langle S_B^x \rangle + 2J_2^\top \langle S_A^x \rangle}{2\omega_A} \tanh(\beta\omega_A) \\
\langle S_A^z \rangle &= \frac{2J_1^\parallel \langle S_B^z \rangle + 2J_2^\parallel \langle S_A^z \rangle + h^z}{2\omega_A} \tanh(\beta\omega_A) \\
\langle S_B^x \rangle &= \frac{2J_1^\top \langle S_A^x \rangle + 2J_2^\top \langle S_B^x \rangle}{2\omega_B} \tanh(\beta\omega_B) \\
\langle S_B^z \rangle &= \frac{2J_1^\parallel \langle S_A^z \rangle + 2J_2^\parallel \langle S_B^z \rangle + h^z}{2\omega_B} \tanh(\beta\omega_B)
\end{aligned} \tag{B4}$$

where

$$\begin{aligned}
\omega_A &= [(2J_1^\top \langle S_B^x \rangle + 2J_2^\top \langle S_A^x \rangle)^2 + \\
&\quad (2J_1^\parallel \langle S_B^z \rangle + 2J_2^\parallel \langle S_A^z \rangle + h^z)^2]^{\frac{1}{2}} \\
\omega_B &= [(2J_1^\top \langle S_A^x \rangle + 2J_2^\top \langle S_B^x \rangle)^2 + \\
&\quad (2J_1^\parallel \langle S_B^z \rangle + 2J_2^\parallel \langle S_A^z \rangle + h^z)^2]^{\frac{1}{2}}
\end{aligned} \tag{B5}$$

In the canted phases where the transversal magnetic fields $\langle S_A^x \rangle$ and $\langle S_B^x \rangle$ are non-zero, the energies ω_A and ω_B can be eliminated from equations (B4) to yield the following important relations:

$$\begin{aligned}
h^z + 2\langle S_A^z \rangle (J_2^\parallel - J_2^\top) + 2\langle S_B^z \rangle J_1^\parallel &= 2J_1^\top \frac{\langle S_B^x \rangle}{\langle S_A^x \rangle} \langle S_A^z \rangle \\
h^z + 2\langle S_B^z \rangle (J_2^\parallel - J_2^\top) + 2\langle S_A^z \rangle J_1^\parallel &= 2J_1^\top \frac{\langle S_A^x \rangle}{\langle S_B^x \rangle} \langle S_B^z \rangle
\end{aligned} \tag{B6}$$

In the limit $h^z \rightarrow \infty$ the Hamiltonian (Equation (B1)) reduces to an effective single operator model:

$$H = -h^z (\langle S_A^z \rangle + \langle S_B^z \rangle) \tag{B7}$$

Consequently for high external fields the system will assume the energetically favorable ferromagnetic phase. In the opposite limit $h^z \rightarrow 0$ and with sufficiently large anti-ferromagnetic

nearest neighbour coupling $J_1^\parallel \ll 0$ the system is dominated by:

$$H = -J_1^\parallel \langle S_A^z \rangle \langle S_B^z \rangle \quad (\text{B8})$$

giving rise to the anti-ferromagnetic state. At medium large fields h^z the two terms Equation (B7) and Equation (B8) balance each other and the transversal ferromagnetic coupling ($J_1^\top > 0$ and $J_2^\top > 0$) become significant, deviating the spins into the x -direction. In regions of higher h^z , where ferromagnetism (Equation (B7)) is more prevalent the system leaps into the canted ferromagnetic phase; for lower h^z it yields the canted anti-ferromagnetic phase. For those sets of coupling constant where all four phases are existent, the corresponding phase transitions are of second order. If, due to choice of constants one or more of those phases, for example the canted anti-ferromagnetic phase, does not appear the resulting canted ferromagnetic to ferromagnetic phase transition is first order.

Now, we may ask the question "For which sets of parameters all four phases appear?". Matsuda and Tsuneto [16] derived relations for all phase transition points at zero temperature. As mentioned in the previously the four phases are distinguished by their order parameters, $m_1 = \langle S_A^x \rangle + \langle S_B^x \rangle$ and $m_2 = \langle S_A^z \rangle - \langle S_B^z \rangle$. Across all second order phase transitions the spin mean-fields $\langle S_x \rangle$ and $\langle S_z \rangle$ are continuous C^0 functions of h^z . Therefore the critical points are determined by Equations (B6), in the limits where the relevant order parameter, m_1 or m_2 , disappears. The canted ferromagnetic phase transits into the ferromagnetic phase when m_1 approaches zero. Hence we set $\langle S_A^x \rangle = \langle S_B^x \rangle \rightarrow 0$ and consequently $\langle S_A^z \rangle = \langle S_B^z \rangle = \frac{1}{2}$. Equation (B6) readily gives:

$$h_{FE-CFE}^z = J_1^\top + J_2^\top - J_1^\parallel - J_2^\parallel \quad (\text{B9})$$

Equally the canted anti-ferromagnetic to anti-ferromagnetic transition is defined by $\langle S_A^z \rangle = -\langle S_B^z \rangle = \frac{1}{2}$ while $\langle S_A^x \rangle \rightarrow 0$ and $\langle S_B^x \rangle \rightarrow 0$. The unknown quotient $\frac{\langle S_A^x \rangle}{\langle S_B^x \rangle}$ is readily eliminated:

$$h_{CAF-AF}^z = \sqrt{(-J_1^\parallel + J_2^\parallel - J_2^\top)^2 - (J_1^\top)^2} \quad (\text{B10})$$

The canted ferromagnetic and the canted anti-ferromagnetic phases coexist where the order parameter of the diagonal long-range order, $m_2 = \langle S_A^z \rangle - \langle S_B^z \rangle$ approaches zero. We replace $\langle S_A^z \rangle$ and $\langle S_B^z \rangle$ in equation (B6) with m_2 and m_1 and retain only linear terms of m_2 . Subtracting and summing up both equations respectively yields:

$$h^z + 2m_2(J_2^\parallel - J_2^\top + J_1^\parallel) = 2J_1^\top m_2$$

$$2m_1(J_2^\parallel - J_2^\top - J_1^\parallel) = -2J_1^\top m_1 \frac{4m_2^2 + 1}{4m_2^2 - 1} \quad (\text{B11})$$

We used that $\sqrt{\langle S_A^z \rangle^2 + \langle S_A^x \rangle^2} = \frac{1}{4}$ at $T=0$. The solution of these two equations determine the critical point which is given by:

$$h_{CFE-CAF}^z = \frac{J_1^\parallel + J_2^\parallel - J_1^\top - J_2^\top}{J_1^\parallel - J_2^\parallel - J_1^\top + J_2^\top} \times \sqrt{(-J_1^\parallel + J_2^\parallel - J_2^\top)^2 - (J_1^\top)^2} \quad (\text{B12})$$

For a particular choice of coupling constants all four phases will exists when:

$$h_{FE-CFE}^z > h_{CFE-CAF}^z > h_{CAF-AF}^z \quad (\text{B13})$$

In other cases, for example where $h_{CFE-CAF}^z < h_{CAF-AF}^z$ the canted anti-ferromagnetic phase is suppressed. The resulting first order canted ferromagnetic - anti-ferromagnetic transition point has to be calculated by finding the state with the lowest internal energy $\langle H \rangle$ and making a comparison between the two.

APPENDIX C: LINK TO MEAN-FIELD SOLUTION

Here we establish a link between the classical mean-field approximation and the random-phase approximation as derived in the previous Chapters. We have already seen that both approximations are mean-field type, involving self consistency equations and that there are two equations (Equation (B6) or Equation (19)) which appear in both approximations. However, the random-phase approximation takes spin-wave/quasi-particle excitations into account whereas the classical mean-field approximation is an effective one operator model exhibiting two energy levels per sub-lattice. Therefore we can now consider if those energy levels, specifically their difference, correspond to certain spin-wave excitations. In this Appendix will show that the classical mean-field approximation is a limiting case of the random-phase approximation in a way in which the integral of the momentum k is restricted to values where:

$$\gamma_1(k) = \gamma_2(k) = 0, \quad (\text{C1})$$

or equally the generalized density of states is:

$$\rho(\gamma_1, \gamma_2) = \delta(\gamma_1)\delta(\gamma_2) \quad (\text{C2})$$




FIG. 4: The rings show the areas where the two lattice generating functions $\gamma_1(k)$ and $\gamma_2(k)$ yield simultaneously zero. If the integral over the first Brillouin zone is restricted to those areas the random-phase approximation devolves into the classical mean-field approximation.

In this limit the self-consistency equation of the random-phase approximation Equation (46) becomes

$$\begin{aligned}
F_{AMF} &= \sum_{\omega_p} [\text{Residue}(\frac{G_A^{yy}(\gamma_1 = 0, \gamma_2 = 0, \omega)}{e^{\beta\omega} - 1}, -\omega_p) \\
&\quad + \text{Residue}(\frac{G_A^{yy}(\gamma_1 = 0, \gamma_2 = 0, \omega)}{e^{\beta\omega} - 1}, -\omega_p)] - \frac{1}{4} = 0 \\
F_{BMF} &= \sum_{\omega_p} [\text{Residue}(\frac{G_B^{yy}(\gamma_1 = 0, \gamma_2 = 0, \omega)}{e^{\beta\omega} - 1}, -\omega_p) \\
&\quad + \text{Residue}(\frac{G_B^{yy}(\gamma_1 = 0, \gamma_2 = 0, \omega)}{e^{\beta\omega} - 1}, -\omega_p)] - \frac{1}{4} = 0
\end{aligned} \tag{C3}$$

Figure 4 shows the wave-vectors k that correspond to $\gamma_1(k) = \gamma_2(k) = 0$ within the first Brillouin zone. Physically, vanishing lattice generating functions ($\gamma_1(k) = \gamma_2(k) = 0$) means that the system loses all information about the lattice structure as it is the case in the classical mean-field approximation where the only information that remains is the number

of nearest and next nearest neighbors. With the lattice generation functions being zero the matrix M of Equation (12) becomes:

$$M^{\text{MF}} = \begin{pmatrix} i\omega & 0 & M_{13} & 0 & 0 & 0 \\ 0 & i\omega & 0 & M_{24} & 0 & 0 \\ -M_{13} & 0 & i\omega & 0 & M_{35} & 0 \\ 0 & -M_{24} & 0 & i\omega & 0 & M_{46} \\ 0 & 0 & M_{53} & 0 & i\omega & 0 \\ 0 & 0 & 0 & M_{64} & 0 & i\omega \end{pmatrix} \quad (\text{C4})$$

where the components are given by:

$$\begin{aligned} M_{13} &= 2h^z + 4\langle S_A^z \rangle J_2^\parallel + 4\langle S_B^z \rangle J_1^\parallel \\ M_{24} &= 2h^z + 4\langle S_B^z \rangle J_2^\parallel + 4\langle S_A^z \rangle J_1^\parallel \\ M_{35} &= 4\langle S_A^x \rangle J_2^\top + 4\langle S_B^x \rangle J_1^\top \\ M_{46} &= -4\langle S_B^x \rangle J_2^\top + 4\langle S_A^x \rangle J_1^\top \\ M_{53} &= -4\langle S_B^x \rangle J_1^\top - 4\langle S_A^x \rangle J_2^\top \\ M_{64} &= -4\langle S_A^x \rangle J_1^\top - 4\langle S_B^x \rangle J_2^\top \end{aligned} \quad (\text{C5})$$

The corresponding Green's functions are given by:

$$\begin{aligned} G_{A_{MF}}^{yy}(\omega) &= \frac{(4J_1^\top \langle S_B^x \rangle + 4J_2^\top \langle S_A^x \rangle) \langle S_A^x \rangle + (4J_1^\parallel \langle S_B^z \rangle + 4J_2^\parallel \langle S_A^z \rangle + 2h^z) \langle S_A^z \rangle}{\omega^2 - (2\omega_1)^2} \\ G_{B_{MF}}^{yy}(\omega) &= \frac{(4J_1^\top \langle S_A^x \rangle + 4J_2^\top \langle S_B^x \rangle) \langle S_B^x \rangle + (4J_1^\parallel \langle S_A^z \rangle + 4J_2^\parallel \langle S_B^z \rangle + 2h^z) \langle S_B^z \rangle}{\omega^2 - (2\omega_2)^2} \end{aligned} \quad (\text{C6})$$

The poles of the Green's functions are given by the eigenvalues of M^{cmf} :

$$\begin{aligned} \omega_1 &= \pm [(2J_1^\top \langle S_B^x \rangle + 2J_2^\top \langle S_A^x \rangle \omega_A)^2 + \\ &\quad (2J_1^\parallel \langle S_B^z \rangle + 2J_2^\parallel \langle S_A^z \rangle + h^z)^2]^{1/2} \\ \omega_2 &= \pm [(2J_1^\top \langle S_A^x \rangle + 2J_2^\top \langle S_B^x \rangle \omega_A)^2 + \\ &\quad (2J_1^\parallel \langle S_B^z \rangle + 2J_2^\parallel \langle S_A^z \rangle + h^z)^2]^{1/2} \end{aligned} \quad (\text{C7})$$

Those energies are, as we have expected, identical to the classical mean field energies given by Equation (B5). The self-consistency equations calculated with Equation (C3) yield:

$$\frac{1}{2\omega_1} [(2J_1^\top \langle S_B^x \rangle + 2J_2^\top \langle S_A^x \rangle) \langle S_A^x \rangle +$$

$$(2J_1^{\parallel}\langle S_B^z\rangle + 2J_2^{\parallel}\langle S_A^z\rangle + h^z)\langle S_A^z\rangle = \frac{\tanh(\beta\omega)}{4} \quad (\text{C8})$$

$$\frac{1}{2\omega_2}[(2J_1^{\top}\langle S_A^x\rangle + 2J_2^{\top}\langle S_{xB}\rangle)\langle S_B^x\rangle + (2J_1^{\parallel}\langle S_A^z\rangle + 2J_2^{\parallel}\langle S_B^z\rangle + h^z)\langle S_B^z\rangle] = \frac{\tanh(\beta\omega)}{4} \quad (\text{C9})$$

Those two equations together with Equation (19) are identically to Equation (B4). Therefore we have shown that the mean-field as derived by Fisher and Liu [6] is indeed a limiting case of the random-phase approximation.

-
- [1] William Butler Yeats, *Last Poems (1936-1939)*, The Municipal Gallery Revisited, st. 7.
 - [2] A. Stoffel and M. Gulácsi, Europhys. Lett. **85**, 20009 (2009).
 - [3] A. J. Stoffel and M. Gulácsi, Eur. Phys. J. **B67**, 169 (2009).
 - [4] A. J. Stoffel and M. Gulácsi, Eur. Phys. J. **B68**, 79 (2009).
 - [5] M. E. Fisher, Rep. Prog. Phys. **30**, 615 (1967).
 - [6] K. -S. Liu and M. E. Fisher, Jour. Low Temp. Phys. **10**, 655 (1973).
 - [7] A. A. Abrikosov, I. P. Gorkov and I. Ye. Dzyaloshinskii, *Methods of Quantum Field Theory in Statistical Physics*, Pergamon Press, 1965.
 - [8] P. E. Bloomfield and N. Nafari, Phys. Rev. **A5**, 806 (1972).
 - [9] N. N. Bogolyubov and S.V. Tyablikov, Doklady Akad. Nauk. S.S.S.R. **126**, 53 (1959).
 - [10] S. V. Tyablikov, Ukrain. Mat. Yhur. **11**, 287 (1959).
 - [11] P. E. Bloomfield and E. B. Brown, Phys. Rev. **B22**, 1353 (1980).
 - [12] A. Auerbach, *Interacting electrons and quantum magnetism*, Springer, 1994.
 - [13] K. W. H. Stevens and G. A. Toombs, Proc. Phys. Soc. **85**, 1307 (1965).
 - [14] T. Matsubara and H. Matsuda, Progr. Theoret. Phys. **16**, 569 (1956).
 - [15] T. Matsubara and H. Matsuda, Progr. Theoret. Phys. **17**, 19 (1957).
 - [16] H. Matsuda, T. Tsuneto, Prog. Theoret. Phys. Suppl. **46**, 411 (1970)